

Isolation and expression analysis of *AGAMOUS*-like genes from *Eucalyptus grandis*

Tomohiko Kato, Takashi Hibino

Forestry Research Institute, Oji Paper Co. Ltd., Kameyama 519-0212, Japan

* E-mail: takashi-hibino@ojipaper.co.jp Tel: +81-5958-5-0122 Fax: +81-5958-5-1528

Received October 28, 2008; accepted December 19, 2008 (Edited by R. Sano)

Abstract In this study, as a first step to achieve a goal to produce genetically-engineered sterile *Eucalyptus*, we focused on the *AGAMOUS*-like genes which are especially involved in stamen formation. Three new distinct cDNA fragments (*EgAGL1*, *EgAGL2* and *EgAGL3*) and two distinct genomic DNA fragments (*EgAGL1* and *EgAGL2*) encoding MADS-box proteins were isolated from *Eucalyptus grandis*. In the present study, three genes have 62–74% homology with the *Arabidopsis AGAMOUS* (AG) gene in deduced amino acid level. *EgAGL1*, *EgAGL2* and *EgAGL3* were classified close to the *AGAMOUS* gene by phylogenetic analysis. *EgAGL1* and *EgAGL2* were strongly expressed in flower buds, and consequently may regulate stamen formation in *Eucalyptus*.

Key words: *AGAMOUS*-like, *Eucalyptus*, MADS-box.

Recently, with the increasing demand for renewable energy, the importance of fiber and chemical materials from woody plants is growing rapidly. In particular, improvement of fast-grow trees such as *Eucalyptus* species, which can grow in drought, cold, or nutrient-poor environments, will be anticipated in the near future world wide (FAO 2004). *Eucalyptus* wood is important as a raw material for industrial pulp and paper making (Higgins 1984; Hillis 1984). The increased bio-mass production from cultivated trees also reduces pressure to exploit native forests. One method to improve the characteristics of cultivated plants is to artificially introduce a certain gene into the target plant using recombinant DNA techniques. However, horizontal spread of the introduced gene derived from a genetically modified (GM) plant by the spread of pollen in the environment is a serious problem (FAO 2004).

Eucalyptus has a distinct floral shape from that of many plants, although the basic floral structures are equivalent to those of most plants (Southerton et al. 1998). In the *Eucalyptus* flower, sepal primordia fuse to each other during the early stage of flower development. Petal primordia are also fused. The fused sepals and petals form a structure called operculum which covers developing reproductive structures before anthesis.

MADS-box genes encode transcription factors that control growth and development, including floral meristem and floral organ formation in plants (Rounsley et al. 1995; Alvarez-Buylla et al. 2000). These proteins

are characterized by the presence of MADS-box domain which is a highly conserved DNA binding domain located in the N-terminal region of the protein (Purugganan et al. 1995). These transcription factors also contain the K-box domain which is involved in protein dimerization. The plant MADS-box genes have been divided into several classes by phylogenetic analysis (Theissen et al. 2000; Parenicova L et al. 2003). The functions of the genes belonging to the same class are thought to be similar across plant species.

In this study, as a first step to achieve a goal to produce genetically-engineered sterile *Eucalyptus*, we focused on the *AGAMOUS*-like genes (Yanofsky et al. 1990; Weigel and Meyerowitz 1994) which are especially involved in reproductive structure, namely stamen and carpel, formation. Although many MADS-box genes have been identified in a number of plant species, few MADS-box genes have been reported in *Eucalyptus* (Kyojuka et al. 1997; Southerton et al. 1998) and they do not have high sequence similarity to *AGAMOUS*. To dissect a molecular basis for floral organ formation in *Eucalyptus*, we isolated *AGAMOUS*-like genes possibly involved in floral organ formation.

Eucalyptus grandis trees were grown in the experimental field at the Forestry Research Institute (Oji Paper Co. Ltd.). RNA was extracted from about 1 g of flower buds from 7-year-old plants using the Plant RNA Isolation Reagent (Invitrogen, USA), according to the supplier's protocol. The SMART kit (Clontech, USA)




Figure 2. Phylogenetic tree of the *Arabidopsis* and rice *AGAMOUS* proteins (AG and OsMADS3), *Arabidopsis* four MADS-box proteins (AP1, AP3, PI and AGL65) and six *Eucalyptus* MADS-box proteins. Amino acid sequences were aligned using the CLUSTALW program and the tree was constructed with the TreeView software.

OsMADS3), *Arabidopsis* four MADS-box genes (*AP1*, *AP3*, *PI* and *AGL65*) and six *Eucalyptus* MADS-box genes using CLUSTALW and the TreeView software. *EgAGL1*, *EgAGL2* and *EgAGL3* were clearly classified close to the *AGAMOUS* gene. Expression of these three genes in flower buds was analyzed by Northern blot analysis (Figure 3). *EgAGL1* and *EgAGL2* mRNAs were strongly expressed in flower buds. Expression of *EgAGL3* in flower buds was weaker than *EgAGL1* and *EgAGL2*. These results indicate that the three *EgAGL* genes are candidate genes for *Eucalyptus AGAMOUS* orthologs.

We examined *EgAGL1* and *EgAGL2* as potential targets for commercial use since the expression of these genes in the flower buds was extremely high. Full-length *EgAGL1* and *EgAGL2* cDNAs were used to screen an *E. camaldulensis* genomic library (Koyama et al. 2006). Two partial genomic DNA sequences with high homology to the *EgAGL1* cDNA and *EgAGL2* cDNA were isolated.

Comparison of the cDNA and genomic sequences of both the *EgAGL1* and *EgAGL2* genes have five introns within the coding region. We isolated upstream sequences (around 1 kbp) from the *EgAGL1* and *EgAGL2* genes, both of which included a TATA-box in a possible promoter region, respectively. Gene accession numbers for the *EgAGL1* and *EgAGL2* promoters are AB465732 and AB465733. These regions will be important components of a transgene to prevent flowering in transgenic *Eucalyptus* trees grown for commercial use.

In the present study, three MADS-box genes were newly identified from flower buds of *Eucalyptus*. These genes have 62–74% homology with the *Arabidopsis AG* gene in deduced amino acid level. *EgAGL1* and *EgAGL2*




Figure 3. Expression analysis of *EgAGL* genes in *E. grandis*. Flower bud mRNA was purified using the PolyATtract mRNA purification system III kit (Promega, USA) according to the supplier's protocol. Purified flower bud mRNA (400 ng) was separated by agarose gel electrophoresis, and then was transferred to a nylon membrane. Blotting and hybridization were performed with Koyama's procedures (Koyama et al, 2006). Expression of *EgAGL1*, *EgAGL2*, and *EgAGL3* was detected using the 3' untranslated region of each gene as a probe. Lane 1, *EgAGL1*; lane 2, *EgAGL2*; lane 3, *EgAGL3*.

were strongly expressed in flower buds, and consequently may regulate stamen and carpel formation in *Eucalyptus*.

Acknowledgements

We are grateful to Ms. Tomoe Shimamura of the Forestry Research Institute of the Oji Paper Co. Ltd for technical assistance with plaque hybridization and sequencing. This work was supported by the New Energy and Industrial Technology Development (NEDO) program, which is part of the "Development of Fundamental Technologies for Controlling the Material Production Process of Plants" project.

References

- Alvarez-Buylla ER, Liljegren SJ, Pelaz S, Gold SE, Burgeff C, Ditta GS, Vergara-Silva F, Yanofsky MF (2000) MADS-box gene evolution beyond flowers: expression in pollen, endosperm, guard cells, roots and trichomes. *Plant J* 24: 457–466
- FAO (2004) *Preliminary review of biotechnology in forestry, including genetic modification. Forest Genetic Resources Working Paper FGR/59E*. Forest Resources Development Service, Forest Resources Division, Rome, pp 1–118
- Higgins HG (1984) Pulp and Paper. In: Hillis WE and Brown AG (eds) *Eucalyptus for wood production*. Academic press Inc, London, pp 290–316
- Hillis WE (1984) Wood quality and utilization. In: Hillis WE and Brown AG (eds) *Eucalyptus for wood production*. Academic press inc, London, pp 259–289
- Koyama T, Kato N, Hibino T, Kawazu T, Kimura T, Sakka K (2006) Isolation and expression analysis of phosphate transporter genes from *Eucalyptus camaldulensis*. *Plant Biotechnol* 23: 215–218
- Kyoizuka J, Harcourt R, Peacock WJ, Dennis ES (1997) *Eucalyptus* has functional equivalents of the *Arabidopsis AP1* gene. *Plant Mol Biol* 35: 573–584

- Parenicova L, de Folter S, Kieffer M, Horner DS, Favalli C, Busscher J, Cook HE, Ingram RM, Kater MM, Davies B, Angenent GC, Colombo L (2003) Molecular and Phylogenetic Analyses of the Complete MADS-Box Transcription Factor Family in Arabidopsis: New Openings to the MADS World. *Plant Cell* 15: 1538–1551
- Purugganan MD, Rounsley SD, Schmidt RJ, Yanofsky MF (1995) Molecular evolution of flower development: diversification of the plant MADS-box regulatory gene family. *Genetics* 140: 345–356
- Rounsley SD, Ditta GS, Yanofsky MF (1995) Diverse Roles for MADS Box Genes in Arabidopsis Development. *Plant Cell* 7: 1259–1269
- Southerton SG, Marshall H, Mouradov A, Teasdale RD (1998) Eucalypt MADS-box genes expressed in developing flowers. *Plant Physiol* 118: 365–372
- Theissen G, Becker A, Di Rosa A, Kanno A, Kim JT, Munster T, Winter K-U, Saedler H (2000) A short history of MADS-box genes in plants. *Plant Mol Biol* 42: 115–149
- Weigel D, Meyerowitz EM (1994) The ABCs of floral homeotic genes. *Cell* 78: 203–209
- Yanofsky MF, Ma H, Bowman JL, Drews GN, Feldmann KA, Meyerowitz EM (1990) The protein encoded by the Arabidopsis homeotic gene *AGAMOUS* resembles transcription factors. *Nature* 346: 35–39